

EVIDENCE FOR A SUPERNOVA IN REANALYZED OPTICAL AND NEAR-INFRARED IMAGES OF GRB 970228

T.J. GALAMA¹, N. TANVIR², P.M. VREESWIJK¹, R.A.M.J. WIJERS³, P.J. GROOT¹, E. ROL¹, J. VAN PARADIJS^{1,4}, C. KOUVELIOTOU^{5,6}, A.S. FRUCHTER⁷, N. MASETTI⁸, H. PEDERSEN⁹, B. MARGON¹⁰, E.W. DEUTSCH¹⁰, M. METZGER¹¹, L. ARMUS¹¹, S. KLOSE¹², B. STECKLUM¹²

Draft version February 5, 2008

ABSTRACT

We present B-, V-, R_c-, I_c-, J-, H-, K- and K'-band observations of the optical transient (OT) associated with GRB 970228, based on a reanalysis of previously used images and unpublished data. In order to minimize calibration differences we have collected and analyzed most of the photometry and consistently determined the magnitude of the OT relative to a set of secondary field stars. We confirm our earlier finding that the early decay of the light curves (before March 6, 1997) was faster than that at intermediate times (between March 6 and April 7, 1997). At late times the light curves resume a fast decay (after April 7, 1997). The early-time observations of GRB 970228 are consistent with relativistic blast-wave models but the intermediate- and late-time observations are hard to understand in this framework. The observations are well explained by an initial power law decay with $\alpha = -1.73^{+0.09}_{-0.12}$ modified at later times by a type-I_c supernova light curve. Together with the evidence for GRB 980326 and GRB 980425 this gives further support for the idea that at least some GRBs are associated with a possibly rare type of supernova.

Subject headings: gamma rays: bursts – supernovae: general

1. INTRODUCTION

The γ -ray burst of February 28, 1997, was the first for which a fading X-ray (Costa et al. 1997) and an optical counterpart (Groot et al. 1997a; van Paradijs et al. 1997) were found. After the counterpart had weakened by several magnitudes, it was found to coincide with an extended object (van Paradijs et al. 1997; Groot et al. 1997b; Metzger et al. 1997a). In subsequent observations with the Hubble Space Telescope (HST) on March 26 and April 7, 1997, it was found that the optical counterpart consisted of a point source and an extended ($\sim 1''$) object (GAL), offset from the point source by $\sim 0.5''$ (Sahu et al. 1997). The extended object is most likely the host galaxy of GRB 970228 and its redshift was recently determined at $z = 0.695 \pm 0.002$ (Djorgovski et al. 1999). The original light curve was compiled from data taken from the literature (Galama et al. 1997). Data obtained until April 1997 suggested a slowing down of the decay of the optical brightness. However, the HST observations of September 1997 (Fruchter et al. 1997, 1999; Castander and Lamb 1998a) showed that after April 1997 the light curve of the point source roughly continued a power-law decay.

We have collected and reanalyzed the photometric information (previously drawn from the literature and presented by

Galama et al. 1997), collected yet unpublished images and present here reassessed Cousins V-, R_c-, and I_c-band optical light curves and spectral flux distributions of the GRB counterpart (Sect. 2.1, 2, 3 and 4). We confirm the result of Galama et al. (1997) that the early decay of the light curves (before March 6, 1997) was faster than that at later times (between March 6 and April 7, 1997) and discuss the observations in terms of the relativistic blast-wave and blast-wave plus underlying supernova models in Sect. 5. We find that the observations are well explained by an initial power law decay with $\alpha = -1.73^{+0.09}_{-0.12}$ modified at later times by a type-I_c supernova light curve. This is in agreement with a recent suggestion by Reichart (1999). Together with similar evidence for GRB 980326 (Bloom et al. 1999) and GRB 980425 (Galama et al. 1998) this gives further support for the idea that at least some GRBs are associated with a possibly rare type of supernova.

2. REANALYSIS OF PREVIOUS OBSERVATIONS

We collected and analyzed most of the photometry of GRB 970228. In Table 1 we provide a log of all the photometry on GRB 970228, obtained in the B, V, R_c, and I_c passbands (corresponding to the Johnson B and Cousins VRI system), some photometry derived from unfiltered passbands and translated to B or R_c, the HST WFPC2 and STIS filters, and

¹Astronomical Institute ‘Anton Pannekoek’, University of Amsterdam, & Center for High Energy Astrophysics, Kruislaan 403, 1098 SJ Amsterdam, The Netherlands

²Department of Physical Sciences, University of Hertfordshire, College Lane, Hatfield, Herts, AL10 9AB, UK

³Department of Physics and Astronomy, SUNY Stony Brook, NY 11794-3800, USA

⁴Physics Department, University of Alabama in Huntsville, Huntsville AL 35899, USA

⁵Universities Space Research Association

⁶NASA/MSFC, Code SD-40, Huntsville AL 35812, USA

⁷Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218, USA

⁸CNR Bologna, Via P. Gobetti 101, 40129 Bologna, Italy

⁹Copenhagen University Observatory, Juliane Maries Vej 30, D-2100, Copenhagen Å, Denmark

¹⁰Astronomy Dept., University of Washington, Box 351580, Seattle WA 98195-1580, USA

¹¹SIRTF Science Center, Caltech, MS 314-6 Pasadena CA 91125

¹²Thüringer Landessternwarte Tautenburg, Sternwarte 5, D-07778, Tautenburg, Germany

the near-infrared J, H, K and K' passbands. In order to minimize calibration differences as a result of use of different filters/instruments and differences in analysis we consistently determined the magnitude of the optical transient (OT) relative to our calibration, which is described in Sect. 2.1, using a set of secondary field stars (Table 2). In Sect. 2.2 and 2.3 we discuss each observation separately.

2.1. Calibration

As part of the ESO Very Large Telescope Unit 1 (VLT-UT1) Science Verification (1998) a series of deep images of the GRB 970228 field were obtained in B, V and R_c (see Table 1). The images were bias-subtracted and flat-fielded in the standard fashion. Through the entire night observations of the Landolt fields SA 99 (around star 253), PG1633+099, MARK A and TPHE (Landolt 1992) were obtained allowing accurate photometric calibration for a number of reference stars (see Table 2). We determined the transformation of the reference stars' magnitude to the Landolt system by fitting for the zero point and the first order extinction and color coefficients.

We also determined a calibration in B, V and R_c from observations of the Landolt fields SA98 (around star 978), PG0918+029 and Rubin 149, taken as part of the Isaac Newton Telescope (INT) observations of March 8 and March 9, 1997 (see Table 1). These two calibrations agree, for stars C-H, to within 0.04, 0.02 and 0.10 magnitudes for B, V and R_c , respectively. The difference between the two calibrations is due to lesser depth of the INT calibration and we estimated that the absolute calibration uncertainty in each filter from the VLT-UT1 observations is 0.05 magnitudes. We used the VLT-UT1 magnitudes for the reference stars C-H and included stars A and B from the INT calibration (stars A and B are not contained in the field of the VLT-UT1 images), correcting for the small zero point offset between the VLT-UT1 and INT calibration. The adopted reference stars' magnitudes are listed in Table 2.

Observations of the field of GRB 970228 in B, V, R_c and I_c (for 240 sec each filter) and of Landolt field SA 98 were obtained March 2, 1999 with the 1-m Johannes Kapteyn Telescope (JKT; La Palma). Of stars A-H only star B was well detected, because of the lesser depth of the JKT images. Therefore, to obtain the I_c -band calibration, we proceeded as follows. First, we determined the magnitudes of star B and six more bright stars in the field of GRB 970228, correcting for the zero point, the atmospheric extinction and also for a first-order color term. We then used the I_c -band calibration of these bright field stars to calibrate stars A-H using the William Herschel Telescope (WHT) March 8, 1997 I_c -band image; these magnitudes are listed in Table 2. For consistency: the calibration in B, V, and R_c for star B is in good agreement with the calibration determined from the VLT-UT1 and INT data (better than 0.07 mag).

Near-infrared images in the J and K band of the field of GRB 970228 and of standard star FS 14 (Casali and Hawarden 1992) were obtained on March 30.2, 1997 UT with the Near Infrared Camera (NIRC) at the Keck-I 10-m telescope. From these observations we obtained a calibration of the nearby star (star H); we found $J = 20.27 \pm 0.10$ and $K = 19.67 \pm 0.10$. Another set of near-infrared images of the field of GRB 970228 and of standard star HD 84800 (Casali and Hawarden 1992) were obtained in J, H and K' with the Calar Alto 3.5-m telescope on March 17.8, 1997 UT. From these observations we found for the nearby star (star H): $J = 20.20 \pm 0.15$, $H = 19.16 \pm$

0.15 and $K' = 19.59 \pm 0.28$. The two calibrations are in good agreement with each other and we adopted their weighted averages: $J = 20.25 \pm 0.10$, $H = 19.16 \pm 0.15$ and $K = 19.66 \pm 0.10$.

2.2. The details of each observation

The earliest image of the OT was reported by Pedichini et al. (1997). The reported magnitude was given relative to the nearby late-type star (LTS; star H in Table 2). This observation was obtained without a filter; we have transformed this unfiltered magnitude to the R_c band, using the reported filter characteristics (Pedichini et al. 1997), taking $V - I_c = 0.50 \pm 0.23$ (from the February 28.99, 1997 UT WHT observations) and assuming a power-law spectrum $F_\nu \propto \nu^\beta$. The resulting value $R_c = 20.0 \pm 0.5$ is consistent with that reported by Pedichini et al. (1997) and Galama et al. (1997).

In the images of Guarnieri et al. (1997)¹³, the OT was blended with the nearby late-type star. We analyzed their February 28.83, 1997 UT R_c -band image and used a large aperture to include both the OT and the late-type star. The resulting magnitude $R_c = 20.30 \pm 0.28$ (OT + GAL + LTS) in our calibration is about 0.8 mag brighter than that reported by Guarnieri et al. (1997); correcting for the LTS we found $R_c = 20.58 \pm 0.28$ (OT + GAL). We have not included the subsequent measurements and upper-limits as given by Guarnieri et al. (1997) as they are consistent with detections of the late-type star only.

Reanalysis of our WHT, INT and New Technology Telescope (NTT; La Silla) images have led to marginal corrections; all corrections, apart from the V band February 28, 1997 (which is 0.3 mag off), are within the errors as presented in Galama et al. (1997). We also present an unpublished R_c -band non-detection (INT March 8.89, 1997 UT).

The March 3.1, 1997 UT observation by Margon et al. (1997) with the Astrophysical Research Consortium's 3.5-m telescope at the Apache Point Observatory was performed using an unfiltered (UF) passband (blue). As the OT was faint in the APO image we determined its position relative to a number of nearby field stars in the February 28.99, 1997 UT V-band WHT image (where the OT is bright) and performed a rotation plus translation to obtain the corresponding pixel position in the APO image; subsequently we have used this position to center the aperture to determine the OT's magnitude (the aperture was chosen such that the extended emission was also contained). The unfiltered OT's magnitude was translated to the B passband by fitting an empirical linear relation between the $UF - B$ magnitude difference and the $B - V$ magnitude difference determined from the reference stars A-H (Table 2). We found $UF - B = 24.03(65) + 0.45(11)(B - V)$ ($\chi^2 = 1.2$ with 3 d.o.f). We obtained a rough estimate of the color of the OT by fitting a temporal power law decay, $F_\nu \propto t^{-\alpha}$, to the V-band data, interpolating the fit to March 9.85, 1997 UT and using the B-band measurement at that same epoch; we found $B - V = 1.10 \pm 0.34$ on March 9.85, 1997 UT. Assuming that the color of the OT did not substantially change this corresponds to $B = 24.5 \pm 0.7$ on March 3.1, 1997 UT. (Note that the error is not dominated by the assumed color). This is about 1.4σ different from the original value (Margon et al. 1997).

Apart from the March 4.9, 1997 UT V-band image (presented by Galama et al. 1997) additional, and yet unpublished, unfiltered (red) and B-band images were obtained with the Nordic Optical Telescope (NOT). The OT was clearly detected in the unfiltered March 4.9, 1997 UT image but not in the V and B-

¹³Note that in Galama et al. 1997 the time of the observation of Guarnieri et al. 1997 is given incorrectly.

band images of March 4.9, 1997 UT nor in the unfiltered image of March 3.9, 1997 UT. Similar to the procedure for the APO data we determined the pixel position of the OT from the February 28.99 V-band WHT image to obtain the magnitude of the OT in the unfiltered NOT March 4.9, 1997 UT image. The unfiltered OT's magnitude was translated to the R_c passband by fitting an empirical linear relation between $UF - R_c$ and $V - R_c$ determined from the reference stars A-I (Table 2). We found $R_c = 22.97(26) + 0.48(16)(V - R_c)$ ($\chi^2 = 4.3$ with 6 d.o.f.). Taking $V - R_c = 1.03$ (corresponding to $V - I_c = 2.24 \pm 0.14$, as observed for the OT with the HST March 26 and April 7, 1997; see below) we found $R_c = 23.46 \pm 0.32$. In a similar way, for $V - I_c = 0.50 \pm 0.23$ (WHT February 28.99, 1997) we obtained $R_c = 23.10 \pm 0.27$; we adopted the mean value $R_c = 23.28 \pm 0.40$. Using the same empirical relation between $UF - R_c$ and $V - R_c$ we obtained $R_c > 22.7 \pm 0.2$ for the unfiltered March 3.9, 1997 UT image (using a conservative $V - I_c = 2.24$).

We also determined the pixel position of the OT to obtain its magnitude in the Keck images of March 6.3 and April 5.8, 1997 UT, using again the February 28.99, 1997 UT V-band WHT image. We found the March 6.3 R_c magnitude, relative to our calibration, to be 0.4 mag brighter than that reported by Metzger et al. (1997a); the April 5.8 magnitude is consistent with that reported by Metzger et al. (1997b).

We have not analyzed the Palomar 200'' September 4, 1997 observation of Djorgovski et al. (1997); it is consistent with detection of the nebulosity only.

We reanalyzed the GRB 970228 HST WFPC2 (F606W and F814W) and STIS images of Fruchter et al. (1999). For the details of the observations we refer the reader to that work. We determined the WFPC2 magnitudes of the OT and stars C, G and H, using a 2 pixel aperture radius. We corrected the instrumental magnitudes for: (i) the 'long versus short anomaly' (Casertano and Mutchler 1998; corrections are -0.16 , -0.24 , -0.12 and -0.19 mag for the WFPC2 observations of March 26 (V), April 7 (V), March 26 (I), April 7 (I), respectively), (ii) the effect of non-optimal Charge Transfer Efficiency (CTE; Whitmore and Heyer, 1997; Whitmore 1998; corrections are ~ -0.06 mag), and (iii) the geometric distortion of WFPC2 images (Holtzman et al. 1995a; corrections are smaller than 0.03 mag). Next we determined the aperture correction from 2 to 11 pixels, used the instrumental zero points from Baggett et al. (1997) and the color corrections from Holtzman et al. (1995b) to obtain the final V- and I_c -band magnitudes. The errors were determined by adding in quadrature the measurement error (Poisson noise), the errors in the above-mentioned corrections, the aperture correction error and a 2% uncertainty for the instrumental zero points. For the OT the error in the magnitude is dominated by the 'long versus short anomaly' correction (the correction is larger for fainter objects). The results for the OT are given in Table 1. We found systematically brighter magnitudes than Fruchter et al. (1999) and Castander and Lamb (1998a), which can be explained by the 'long versus short anomaly' correction that we applied. We found the OT to be redder ($V - I_c = 2.22 \pm 0.25$ and $V - I_c = 2.26 \pm 0.30$, March 26 and April 7, 1997 UT, respectively) than found by Fruchter et al. (1999), but consistent with the result of Castander and Lamb (1998a). For stars C, G and H we obtained (V, I_c) = (21.76 ± 0.05 , 20.10 ± 0.05), (23.46 ± 0.05 , 22.52 ± 0.07) and (22.85 ± 0.04 , 21.04 ± 0.04), respectively. Comparing this with the ground-based calibration

(see Table 2) we found the agreement for the bright star C to be excellent (better than 0.02 mag), for star H we found 0.15 and 0.12 mag differences and the V-band HST magnitude of star G was 0.22 mag off.

The HST STIS observations were taken in unfiltered mode, meaning that the response covers a wide wavelength range from about 2000 to 10000 Å. To convert the observed counts into broadband magnitudes we had to assume therefore something about the spectral energy distribution. Our procedure was as follows: (i) measure the counts in a 2-pixel radius aperture; (ii) add an aperture correction of -1.15 mags based on brighter stars in the field, assuming that 10% of the light falls outside a large 11 pixel radius aperture; (iii) assume a power law slope with spectral index chosen to be consistent with the earlier WFPC2 observations ($\beta = -4.7$), and use the SYNPHOT routine CALCPHOT to determine the corresponding broad-band magnitudes to explain the observed counts. The result, corrected for the small 12% renormalization found by Landsman (1997), was $V = 28.80$, $R_c = 27.69$ and $I_c = 26.54$. Although the uncertainty in the measured counts is only 10%, the considerable dependence on the shape of the assumed spectrum, and the difficulty of absolutely calibrating the wide STIS passband (see below), leads us to adopt an uncertainty of 0.4 magnitudes for each band.

To check the STIS result we tried an alternative calibration using our WFPC2 photometry for the two brightest field stars, but still relying on SYNPHOT to calculate the appropriate colour correction, and found a STIS magnitude of $V = 28.58$. This discrepancy is larger than the formal errors, and is probably indicative of the difficulty of precisely calibrating the very wide unfiltered STIS CCD passband.

We collected and rereduced the near-infrared J- and K-band images of Soifer et al. (1997), obtained on March 30.2 and 31.2, 1997 UT with NIRC on the Keck-I 10-m telescope, and the J-, H- and K'-band images of Klose, Stecklum and Tuffs (1997), obtained with the Calar Alto 3.5-m telescope on March 17.8, 1997 UT using the near-infrared camera MAGIC. The infrared frames were reduced by first removing bad pixels and combining at least five frames around each object image to obtain a sky image. This sky image was then subtracted after scaling it to the object image level.

A source was detected in both bands J and K in the Keck-I observations. Relative to the nearby star (star H) we found $J = 23.27 \pm 0.15$ (OT + GAL; sum of the March 30.3 and March 31.2, 1997 UT data) and $K = 22.85 \pm 0.25$ (March 30.2, 1997 UT) within a large $0''.75$ aperture radius. The J-band magnitude is consistent with the earlier determination (Soifer et al. 1997), but our K-band determination is 0.8 mag fainter. Fruchter et al. (1999) detected the host galaxy at $K = 22.8 \pm 0.3$ and so our result is consistent with detection of the host galaxy only.

We did not detect the source in the Calar Alto J, H and K' images and determined the following limiting magnitudes: $J > 21.45$, $H > 20.02$ and $K' > 20.13$ (3σ). We confirm the non detections in H and K' by Klose et al. (1997), but do not confirm their detection of the source in J¹⁴.

2.3. The host galaxy

The host galaxy of GRB 970228 is clearly detected in the HST WFPC2 images, but at low signal-to-noise. Working with cosmic-ray and hot-pixel cleaned images, we first subtracted

¹⁴The J-band magnitude reported by Klose et al. (1997) is affected by a transient warm pixel at the position of the source that was not obvious in the bad pixel mask.

point-spread functions positioned and scaled to remove the image of the OT from each. For the two F606W images we then performed aperture photometry in apertures of increasing size, taking a sky estimate from the surrounding area which was free of other objects. We found that the counts reached an asymptote, within the noise, by a radius of 25 pixels ($\sim 1''.1$). We calibrated this using the procedure of Holtzman (1995b), finding an average $V_{606} = 25.5 \pm 0.3$, where the error is dominated by the photon and read noise.

In the F814W image the host galaxy is even fainter, and we estimated its color by comparing the counts within a 7 pixel radius aperture with the counts in the same aperture on the F606W image. Thus we found a color $V_{606} - I_{814} \sim 1.1$ which is, correcting for the Galactic extinction, reasonable for a galaxy at this magnitude (Smail et al. 1995). This could then be used to estimate color corrections to the standard systems, and we concluded $V = 25.75 \pm 0.35$ and $I_c = 24.7 \pm 0.5$.

The host galaxy is much better detected on the STIS image, although we must assume a spectral shape in order to estimate broad-band magnitudes. Using CALCPHOT we found a good match to the WFPC2 colours with a suitably redshifted and reddened S_c galaxy spectrum¹⁵. Based on this, we infer $V = 25.77$, $R_c = 25.22$ and $I_c = 24.73$ from the STIS data, which is in good agreement with the other determinations. We adopt an error of 0.2 mag to account for the systematic uncertainty introduced by the necessity of assuming the spectral shape.

We did not detect the host galaxy of GRB 970228 in the ESO VLT-UT1 Science Verification images and determined the 3σ limit for detection using an aperture radius of 1.5 times the seeing, which corresponds to $\sim 1''.5$. The results are shown in Table 1. The V and R_c -band upper limits for the host galaxy show that the host galaxy is fainter than the earlier determination (Sahu et al. 1997), but consistent with the result of Fruchter et al. (1999), Castander and Lamb (1998a) and with our estimate from the WFPC2 images.

3. THE LIGHT CURVES

In the interpolations to the R_c band (see Fig. 1) we have assumed that the spectra of both the point source and the host galaxy, are smooth (i.e., not dominated by emission lines). We have used the relation between the color indices $V - R_c$ and $V - I_c$ given by Thé et al. (1984) for late-type stars; for bluer stars we have inferred this relation from the tables given by Johnson (1966) for main-sequence stars and the color transformations to the Cousins VRI system given by Bessel (1976). We have tested the validity of these color-color relations from numerical integrations of power law flux distributions and of Planck functions, and conclude that if the flux distributions of the OT and the host galaxy are smooth, the uncertainty in the interpolated R_c magnitude is unlikely to exceed 0.1 mag (Galama et al. 1997). For the HST September 4, 1997 observation we have assumed that the colors of the OT remained constant during the late-time decay (i.e., taking the observed $V - I_c = 2.24$ from the HST March 26 and April 7, 1997 observations; see Sect. 2.2). Finally, we corrected for the contribution of the host galaxy emission ($V = 25.77 \pm 0.20$, $R_c = 25.22 \pm 0.20$ and $I_c = 24.73 \pm 0.20$) and obtained the V-, R_c -, and I_c -band light curves of the OT (Fig. 1).

We have fitted power laws, $F \propto t^\alpha$, to the light curves (ignoring upper limits) and found: $F_V = (11.7 \pm 1.0)t_d^{-1.32 \pm 0.04} \mu\text{Jy}$ ($\chi^2 = 0.2$ with 2 d.o.f), $F_{R_c} = 12.7^{+1.2}_{-1.8}t_d^{-1.14 \pm 0.04} \mu\text{Jy}$ ($\chi^2 = 9.7$ with 8

d.o.f), and $F_{I_c} = 15.2^{+3.1}_{-2.5}t_d^{-0.92 \pm 0.06} \mu\text{Jy}$ ($\chi^2 = 4.5$ with 2 d.o.f), where t_d is the time in days. These fits are indicated with dotted lines in Fig. 1. We also fitted a power law to the R_c -band light curve at intermediate times (between 3 and 50 days) and found $F_{R_c} = 3.4^{+2.4}_{-1.4}t_d^{-0.73 \pm 0.17} \mu\text{Jy}$ ($\chi^2 = 1.3$ with 4 d.o.f). The fit is indicated with a thick line in Fig. 1. For the late-time detections (after 20 days) we found: $F_V = (11.7 \pm 1.3)t_d^{-1.14 \pm 0.21} \mu\text{Jy}$ ($\chi^2 = 0.2$ with 1 d.o.f), $F_{R_c} = 21.5^{+23}_{-11}t_d^{-1.27 \pm 0.20} \mu\text{Jy}$ ($\chi^2 = 0.3$ with 1 d.o.f) and $F_{I_c} = 50^{+48}_{-24}t_d^{-1.25 \pm 0.19} \mu\text{Jy}$ ($\chi^2 = 0.9$ with 1 d.o.f), where t_d is the time in days. These fits are indicated with dashed lines in Fig. 1.

4. OPTICAL/NEAR-INFRARED TO X-RAY SPECTRA

For three epochs we have reconstructed the spectral flux distribution of the OT, corresponding to: (i) 1997 February 28.99 UT (V, R_c and I_c data from Table 1 and 2-10 keV flux from Costa et al. 1997), (ii) March 9.89 UT (R_c band measurement from Table 1 and ROSAT 0.1-2.4 keV unabsorbed flux from Frontera et al. 1998), and (iii) March 30.8 UT (See Fig. 2). We subtracted the host galaxy flux from ground-based measurements, corrected the OT fluxes for Galactic foreground absorption, $A_V = 0.78 \pm 0.12$ (from Schlegel, Finkbeiner and Davis 1998; but see also the discussion by González, Fruchter, and Dirsch 1999, and Castander and Lamb 1998b), and brought measurements to the same epoch using the power-law fits of Sect. 3. We note that the spectral flux distribution on March 30.8 UT (Fig. 2) is different from that of Reichart (1999) because in our reanalysis the optical transient was not detected in the K band. The resulting spectral flux distribution therefore supersedes that of Reichart (1999) who obtained the K-band magnitudes of the OT and the host from the literature.

We fitted the optical to X-ray spectral flux distribution of the first epoch (February 28.99, 1997 UT) with a power law and an exponential optical extinction law, $F_\nu \propto \nu^\beta e^{-\tau}$, where we assumed that the extinction optical depth, $\tau \propto \nu$. The data did not allow determination of the host galaxy extinction; the fit provided a negative extinction ($A_V < 0.50$; 90 % confidence) and was subsequently fixed at zero. We found a spectral slope, $\beta = -0.780 \pm 0.022$ (reduced χ -squared, $\chi^2_r = 4/2$). For comparison, the X-ray power-law photon index observed ~ 12 hours after the burst (2-10 keV; Costa et al. 1997) corresponds to a spectral slope $\beta = -1.1 \pm 0.3$.

5. DISCUSSION

Most X-ray and optical/infrared afterglows display a power law temporal decay; this is a prediction for relativistic blast-wave models of GRB afterglows, in which a relativistically expanding shock front, caused by an energetic explosion in a central compact region, sweeps up the surrounding medium and accelerates electrons in a strong synchrotron emitting shock (e.g., Mészáros and Rees 1997). Assuming a single power law and ignoring upper limits for the moment, the V-, R_c - and I_c -band light curves (Fig. 1) indicate that the power law decay rate is an increasing function of frequency; for the V, R_c and I_c band the power-law decay index $\alpha = -1.32 \pm 0.04$, -1.14 ± 0.04 , and -0.92 ± 0.06 , respectively. This implies that as a function of time the OT becomes redder and redder. Such a relation has not been observed previously, and is not what is expected in standard relativistic blast-wave models, where the power law index is, to first order, independent of frequency.

¹⁵From the Kinney-Calzetti atlas, see the Synphot User Guide 1998, Space Telescope Science Institute.

Taking the upper limits into account, we find that the March 4.87, 1997 UT NOT V-band non-detection is significantly below the fit. Also, the R_c -band light curve suggests that the optical emission is weaker around that time (between 3 and 10 days) than expected from a single power law. The V and R_c data suggest that the initial decline was faster than the decline at intermediate times. From the V-band February 28.99 and March 4.87, 1997 UT data we find that the early-time decay was steeper than $\alpha = -1.4$ (3σ), while at intermediate times (around ~ 10 days) it was $\alpha = -0.73 \pm 0.17$ (R_c). This structure in the light curve was recognized early on by Galama et al. (1997), who showed that the early decay of the light curves (before March 6, 1997) was faster than that at later times (between March 6 and April 7, 1997). The earlier analysis of the March 26 and April 7, 1997 HST WFPC2 images yielded brighter estimates of the host galaxy magnitudes (Sahu et al. 1997) than our reanalysis (see also Fruchter et al. 1999; Castander and Lamb 1998a). Because of this, the correction to the ground-based data for the host galaxy emission applied in this work is smaller than that of Galama et al. (1997), and thereby the effect is somewhat reduced, but it is still present (see also Reichart 1999) and consistent with the earlier result.

The late-time light curves show that the rate of decay increased again after ~ 35 days. The late-time decays in V, R_c and I_c are consistent with being frequency independent (i.e., similar decays in V, R_c and I_c), with a power-law index $\alpha = -1.2$. The back extrapolation of the late-time fit in the V-band predicts quite well the earliest V-band data, but this is not the case for the R_c - and I_c -band light curves. We conclude that a single power-law is not a good representation of the light curves; the overall shape of the lightcurves is that of an initial fast decay (before $\sim$ March 6, 1997) followed by a ‘plateau’ (slow decay; between $\sim$ March 6 and $\sim$ April 7, 1997) and finally the light curves resume a fast decay (after $\sim$ April 7, 1997).

For synchrotron radiation, by a power-law energy distribution of electrons, at low frequencies, and for an adiabatic evolution of the blast wave we can distinguish the following two cases, each of which has its own relation between the spectral slope β and the power law decay index, α (Sari, Piran and Narayan 1998): (i) both the peak frequency ν_m and the cooling frequency ν_c are below the optical waveband. Then $\alpha = 3\beta/2 + 1/2 = -0.67 \pm 0.03$ (taking the measured spectral slope, $\beta = -0.78$ on February 28.99, 1997 UT) (ii) ν_m has passed the optical waveband, but ν_c has not yet. In that case $\alpha = 3\beta/2 = -1.17 \pm 0.03$. The temporal decay values measured at early times are in reasonable agreement with case (ii), but certainly not with case (i), suggesting that the peak frequency ν_m is below the optical passband and that the cooling frequency ν_c is near or above the X-ray (2-10 keV) passband; such a high cooling frequency appears to be the fairly general case in GRB afterglows (e.g., Bloom et al. 1998; Vreeswijk et al. 1999; Galama et al. 1999). The fit to the optical to X-ray spectrum of February 28.99, 1997 UT suggests that extinction at the host is not very significant ($A_V < 0.5$; corresponding to a restframe extinction $A_V < 0.3$). At the second epoch we measured a spectral slope, $\beta = -0.52 \pm 0.08$, on March 9.89, 1997 UT. For case (ii) this would correspond to a power-law decay of $\alpha = -0.78 \pm 0.12$, consistent with a shallower decay at intermediate times. We thus find that the observations of the early-time

afterglow are roughly consistent with the predictions of simple relativistic blast-wave models (see also Wijers, Rees and Mészáros 1997; Waxman 1997; Reichart 1997), except that we find evidence for a flattening of the decay rate around ~ 10 days.

As mentioned above, the frequency dependence of the rate of decay of the light curves corresponds to a reddening of the optical afterglow with time ($V - I_c = 0.50 \pm 0.23$ at February 28.99, 1997 UT and $V - I_c = 2.24 \pm 0.14$ at March 26 and April 7, 1997). We are confident about this large change in color as the HST calibration of three field stars agrees well with that of the ground-based calibration. Assuming a power-law spectrum, the optical spectral slopes (Galactic extinction corrected) from the intermediate-time V and I_c HST observations are $\beta = -3.8 \pm 0.7$ (March 26, 1997) and $\beta = -4.0 \pm 0.8$ (April 7, 1997). These steep spectra cannot be explained by extinction at the host galaxy; we find a staggering $A_V = 4.5 \pm 1.0$ (fitting as before for an exponential optical extinction law, $F_\nu \propto \nu^\beta e^{-\tau}$, with $\tau \propto \nu$, and fixing the spectral slope at $\beta = -0.78$), which would imply that the extinction increased with time from negligible ($A_V < 0.5$) to very significant ($A_V = 4.5 \pm 1.0$); this seems quite unlikely. A steepening of the spectral slope may be expected from a number of spectral transitions: e.g., the passage of the cooling frequency or the transition to a Sedov-Taylor non-relativistic expansion (e.g., Sari et al. 1998; Waxman, Kulkarni and Frail 1998). However, such spectral transitions will be accompanied with an increase in the rate of decay of the light, which we do not observe here. The discrepancy is even larger if one considers the spectral flux distribution around the third epoch (March 30.8, 1997 UT; Fig. 2); the spectrum shows an unexpected turnover around 3×10^{14} Hz (Fruchter et al. 1999). This break plus the very red spectrum in V and I_c in the intermediate-time optical flux distribution is not consistent with the predictions of simple relativistic blast-wave models (see also Reichart 1999; Fruchter et al. 1999).

What then could explain these observations? Bloom et al. (1999) observed a similarly red spectrum for the late-time emission from GRB 980326. The authors argue that this is due to a supernova (SN) that dominated the light at late times. Reichart (1999) suggests that this is also the case for GRB 970228. In this hypothesis the early light curve is dominated by a (power-law decaying) GRB afterglow (produced by the relativistic blast wave), while at late times the light curve contains a significant contribution from an underlying supernova. Such would be the natural outcome of the ‘collapsar/microquasar’ models in which GRBs arise in jets that are formed in the core collapse to a black hole of massive stars (Woosley 1993; Paczyński 1998). Such massive stars will, prior to collapse, have lost their hydrogen envelopes and so we expect the supernova to be of type I_b/c (Woosley 1993; Woosley, Eastman and Schmidt 1999; Bloom et al. 1999).

Following Bloom et al. (1999) and Reichart (1999) we constructed simulated supernova light curves and spectral flux distributions as follows. First we combined three sets of Galactic extinction corrected U-, B-, V-, R_c - and I_c -band observations of the well-observed, unusual type- I_c SN 1998bw (Galama et al. 1998; McKenzie and Schaefer 1999; Vreeswijk et al. 1999), which is likely to be associated with GRB 980425 (Galama et al. 1998; Kulkarni et al. 1998)¹⁶. We then redshifted the

¹⁶The association between SN 1998bw and GRB 980425 has been considered uncertain because the behavior of a another source in the error box of GRB 980425, the fading X-ray source 1SAX J1935.3-5252, resembled that of a typical GRB afterglow. However, reanalysis of NFI BeppoSAX observations of the GRB 980425 field by Pian et al. (1999) shows that its behavior is unlike that of previously observed GRB afterglows. SN 1998bw remains not typical for GRB afterglows because of its low redshift, $z = 0.0085$.

lightcurves to the redshift of GRB 970228, $z = 0.695$ (Djorgovski et al. 1999), which consists of wavelength shifting, time-profile stretching, and dimming (we assumed a cosmology with $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$ and $\Omega_0 = 0.3$). To obtain the corresponding redshifted light curves in the V, R_c and I_c bands we assumed a power-law spectral shape to interpolate (and extrapolate) the redshifted SN 1998bw light curves to the central wavelengths of the V, R_c and I_c bands. (Note that the redshifted central wavelength of the U band is near V and R_c , and for the B band is near I_c .) Finally we corrected the SN light curves for the Galactic extinction in the direction of GRB 970228 ($A_V = 0.78 \pm 0.12$). These light curves are shown in Fig. 3.

The supernova spectral flux distribution corresponding to the epoch of March 30.8, 1997 UT (30.7 days after GRB 970228) is shown in Fig. 2 (here we did the opposite: we corrected the GRB 970228 spectral flux distribution for Galactic extinction). The resemblance of the GRB 970228 and the redshifted SN 1998bw spectral flux distributions is remarkable (see also Reichart 1999).

At early times the R_c -band light curve (see Fig. 3) will not be very much dominated by SN emission. If we attribute the early part ($t < 8$ days) of the R_c -band light curve to GRB afterglow and fit a power-law temporal decay to the data we find a temporal decay index $\alpha = -1.46 \pm 0.16$ ($\chi^2 = 1.5$ with 3 d.o.f.). This value of the decay constant is consistent with that in X rays, $\alpha = -1.33^{+0.11}_{-0.13}$ during the first ~ 4 days (2-10 keV; Costa et al. 1997) and $\alpha = -1.50^{+0.35}_{-0.23}$ up to ~ 10 days (0.1-2.4 keV; Frontera et al. 1998). It is also consistent with the earlier determination (Galama et al. 1997; see also Reichart 1999).

Next we took a single value for the power-law decay in V, R_c and I_c , computed the sums of these power-law decays plus supernova light curves, and fitted for the power-law decay index, α , by minimizing the χ -squared of the V, R_c , I_c light curves and the resulting fit. But first, the assumption of a constant color that was used to derive a V-band magnitude from the September 1997 unfiltered STIS observation is no longer valid as in this hypothesis at late times the SN emission will dominate the light and so we expect the light to have reddened (see Fig. 3). Thus, in order to convert the observed counts into broadband magnitudes we assumed that the spectral energy distribution was similar to that of SN 1998bw, redshifted to $z = 0.695$. Our procedure was as follows: (i) we used the redshifted U, B, V, R_c , I_c spectral-flux distribution of SN 1998bw (on day 111, corresponding to the STIS epoch at 188 days), (ii) we corrected for Galactic extinction, $A_V = 0.78 \pm 0.12$, interpolating to the redshifted central wavelengths of the U, B, V, R_c , I_c bands using the Galactic extinction curve of Cardelli, Clayton and Mathis (1989) to obtain the spectral flux distribution as it would be observed in the direction of GRB 970228, and (iii)

we used this spectral flux distribution in the SYNPHOT routine CALCPHOT, and applied the Landsman (1997) correction, to determine corresponding broad-band magnitudes, $V = 28.56$, $R_c = 27.42$ and $I_c = 26.43$. Again, we adopt an uncertainty of 0.4 mag to make some allowance for the sensitivity of the result to the assumed spectral shape. Finally we minimized the χ -squared and found $\alpha = -1.73^{+0.09}_{-0.12}$ ($\chi^2 = 17.5$ with 14 d.o.f.; we note that the χ -squared is not strictly correct since a few of the data points are not independent). The fit is excellent in view of the uncertainties in the model. Similar results were presented by Reichart (1999). The fit is somewhat better than that presented by Reichart (1999), presumably because of our consistent reanalysis of all the original images, and, in particular, because our STIS magnitudes were calculated using the knowledge of the spectral shape of the template supernova.

The model has very few assumptions: the power law index α is the only free parameter and we did not allow the peak luminosity of the SN to be a free parameter but used the SN 1998bw observations as they are. SN 1998bw is, of course, an unusual example of the class, being very bright and rapidly expanding, but its coincidence with GRB 980425 makes it the only template we can reasonably choose for now. Also, from the ‘collapsar’ model we expect a GRB to be accompanied by a supernova of type I_b/c . In any case, the light curves (Fig. 3) as well as the intermediate-time spectral flux distribution (Fig. 2) and intermediate-time colors are remarkably well explained by the sum of a power-law plus supernova light curve. It can explain the apparent transition from a rapid to a slow and again a rapid decline of the light curves, is in good agreement with the upper limits in the light curves and it can account for the unusual intermediate-time red color $V - I_c = 2.24$ and the peaked spectral flux distribution on March 30.8, 1997 UT (see also Reichart 1999). These features cannot be accounted for by simple relativistic blast-wave models. We thus find the data to be in agreement with the hypothesis of Reichart (1999) that a supernova dominated the light at late times. Together with the evidence for GRB 980326 (Bloom et al. 1999) and GRB 980425 (Galama et al. 1998) this is further support for the idea that at least some GRBs are associated with a possibly rare type of supernova.

We wish to thank Drs. Tufts, Soifer, Neugebauer, Guarnieri, Bartolini and Piccioni for observations of GRB 970228. This work is partly based on observations collected at the European Southern Observatory, Paranal, Chile (VLT-UT1 Science Verification Program). T.J. Galama is supported through a grant by NFRA under contract 781.76.011. PMV and ER are supported by the NWO Spinoza grant. CK acknowledges support from NASA grant NAG 5-2560.

REFERENCES

- Baggett, S., Casertano, S., Gonzaga, S., Ritchie, C. 1997, Instrument Science Report WFPC2 97-10
 Bessel, M.S. 1976, PASP, 88, 557
 Bloom, J.S. et al. 1998, ApJ, 508, L21
 Cardelli, J.A., Clayton, G.C. and Mathis, J.S. 1989, ApJ, 345, 245
 Casali, M.M. and Hawarden, T.G. 1992, JCMT-UKIRT Newsletter, No. 3, 33
 Casertano, S. and Mutchler, M. 1998, Instrument Science Report WFPC2 98-02
 Castander, F.J. and Lamb, D.Q. 1998a, in press, ApJ, astro-ph/9807195
 Castander, F.J. and Lamb, D.Q. 1998b, in press, ApJ, astro-ph/9807194
 Costa, E. et al. 1997, Nature, 387, 783
 Djorgovski, S.G., et al. 1997, IAU Circ., 6732
 Djorgovski, S.G. et al. 1999, GCN Circular, No. 289
 Frontera, F. et al. 1998, A&A, 334, 69
 Fruchter, A.S. et al. 1997, IAU Circ., 6747
 Fruchter, A.S. et al. 1999, ApJ, 516, 683
 Galama, T.J. et al. 1997, Nature, 387, 479
 Galama, T.J. et al. 1998, Nature, 395, 670
 Galama, T.J. et al. 1999, Nature, 398, 394
 González, R.A., Fruchter, A.S., and Dirsch, B. 1999, ApJ, 515, 69
 Groot, P.J. et al. 1997a, IAU Circ., 6584
 Groot, P.J. et al. 1997b, IAU Circ., 6588
 Guarnieri, A. et al. 1997, A&A, 328, L13
 Holtzman, J. et al. 1995a, PASP, 107, 156
 Holtzman, J.A., Burrows, C.J., Casertano, S., Hester, J.J., Trauger, J.T., Watson, A.M., Worthey, G., 1995b, PASP, 107, 1065
 Johnson, H.L. 1966, ARA&A, 4, 191
 Klose, S., Stecklum, B., Tufts, R. 1997, IAU Circ., 6611
 Kulkarni, S.R. et al. 1998, Nature, 395, 663

- Landolt, A.U. 1992, AJ, 104, 340
 Landsman, W. 1997, STIS CCD image mode sensitivity, Internal STIS Calibration Document.
 Margon, B., Deutsch, E.W., Lamb, D.Q., Castander, F.J. 1997, IAU Circ., 6618
 McKenzie, E.H. and Schaefer, B.E. 1999
 Mészáros, P. and Rees, M.J. 1997, ApJ, 476, 232
 Metzger, M.R. et al. 1997a, IAU Circ., 6588
 Metzger, M.R. et al. 1997b, IAU Circ., 6631
 Paczyński, B. 1998, ApJ, 494, L45
 Pedichini et al. 1997, A&A, 327, L32
 Pian, E. et al. 1999, A&AS, in press
 Reichart, D.E. 1997, ApJ, 485, L57
 Reichart, D.E. 1999, ApJ, in press, astro-ph/9906079
 Sahu, K. et al. 1997, Nature, 387, 476
 Sari, R., Piran, T., and Narayan, R. 1998, ApJ, 497, L17
 Schlegel, D.J., Finkbeiner, D.P. and Davis, M. 1998, ApJ500, 525
 Soifer, B. et al. 1997, IAU Circ., 6619
 Smail, I., Hogg, D.W., Yan, L. and Cohen, J.G. 1995, ApJ, 449, L105
 Thé, P.S., Steenman, H., Alcaïno, G., 1984, A&A, 132, 385
 Van Paradijs, J. et al. 1997, Nature, 386, 686
 Vreeswijk, P.M. et al. 1999, in press, ApJ
 Waxman, E. 1997, ApJ, 485, L5
 Waxman, E., Kulkarni, S.R. and Frail, D.A. 1998, ApJ, 497, 288
 Whitmore, B. and Heyer, I. 1997, Instrument Science Report WFPC2 97-08
 Whitmore, B. 1998, Technical Instrument Report WFPC2 98-01
 Wijers, R.A.M.J., Rees, M.J., and Mészáros, P. 1997, MNRAS, 288, L51
 Woosley, S.E. 1993, ApJ, 405, 273
 Woosley, S.E., Eastman, R.G. and Schmidt, B.P. 1999, ApJ, 516, 788

TABLE 1

Summary of optical observations. Upper limits are three sigma. Abbreviations: OT, optical transient; GAL, host galaxy; RAO, Rome Astrophysical Observatory; BUT, Bologna University Telescope; WHT, William Hershell Telescope; APO, Apache Point Observatory; NOT, Nordic Optical Telescope; INT, Isaac Newton Telescope; CA, Calar Alto 3.5-m telescope; HST, Hubble Space Telescope; P5m, Palomar 200-inch; VLT-UT1, Very Large Telescope Unit 1.

Date (UT)	Telescope	Exp. time (sec)	Magnitude	Remarks
Feb. 28.81 1997	RAO		$R_c = 20.0 \pm 0.5$	Unfiltered OT+GAL
Feb. 28.83 1997	BUT		$R_c = 20.58 \pm 0.28$	OT+GAL
Feb. 28.99 1997	WHT		$V = 21.01 \pm 0.10$	OT+GAL
Feb. 28.99 1997	WHT		$I_c = 20.51 \pm 0.21$	OT+GAL
Mar. 03.10 1997	APO		$B_j = 24.5 \pm 0.7$	OT+GAL
Mar. 03.90 1997	NOT		$R_c > 22.70 \pm 0.20$	Unfiltered OT+GAL
Mar. 04.86 1997	NOT		$R_c = 23.28 \pm 0.40$	Unfiltered OT+GAL
Mar. 04.87 1997	NOT		$V > 23.96 \pm 0.12$	OT+GAL
Mar. 04.89 1997	NOT		$B > 23.87 \pm 0.10$	OT+GAL
Mar. 06.32 1997	Keck		$R_c = 23.60 \pm 0.22$	OT+GAL
Mar. 08.86 1997	INT		$V > 24.00 \pm 0.10$	OT+GAL
Mar. 08.88 1997	WHT		$I_c > 22.68 \pm 0.10$	OT+GAL
Mar. 08.89 1997	INT		$R_c > 23.35 \pm 0.08$	OT+GAL
Mar. 09.85 1997	INT		$B = 25.43 \pm 0.31$	OT+GAL
Mar. 09.89 1997	INT		$R_c = 23.75 \pm 0.28$	OT+GAL
Mar. 13.00 1997	NTT		$R_c = 24.36 \pm 0.15$	OT+GAL
Mar. 17.84 1997	CA		$J > 21.45 \pm 0.15$	OT+GAL
Mar. 17.83 1997	CA		$H > 20.02 \pm 0.15$	OT+GAL
Mar. 17.81 1997	CA		$K' > 20.13 \pm 0.15$	OT+GAL
Mar. 26.38 1997	HST		$V = 25.98 \pm 0.22$	OT
Mar. 26.47 1997	HST		$I_c = 23.76 \pm 0.11$	OT
Mar. 30.27 1997	Keck		$K = 22.85 \pm 0.25$	OT+GAL
Mar. 30.30 + 31.26 1997	Keck		$J = 23.27 \pm 0.15$	OT+GAL
Apr. 05.76 1997	Keck		$R_c = 24.97 \pm 0.25$	OT+GAL
Apr. 07.22 1997	HST		$V = 26.36 \pm 0.26$	OT
Apr. 07.30 1997	HST		$I_c = 24.10 \pm 0.15$	OT
Mar. 26 + Apr. 7 1997	HST		$V = 25.75 \pm 0.35$	GAL
Mar. 26 + Apr. 7 1997	HST		$I_c = 24.7 \pm 0.5$	GAL
Sep. 04 1997	P5m		$R = 25.5 \pm 0.5$	GAL
Sep. 04.71 1997	HST		$V = 28.80 \pm 0.40$	OT
Sep. 04.71 1997	HST		$R_c = 27.69 \pm 0.40$	OT
Sep. 04.71 1997	HST		$I_c = 26.54 \pm 0.40$	OT
Sep. 04.71 1997	HST		$V = 25.77 \pm 0.20$	GAL
Sep. 04.71 1997	HST		$R_c = 25.22 \pm 0.20$	GAL
Sep. 04.71 1997	HST		$I_c = 24.73 \pm 0.20$	GAL
Sep. 01.35 1998	VLT-UT1	1800	$B > 26.08 \pm 0.08$	GAL
Sep. 01.35 1998	VLT-UT1	1200	$V > 25.33 \pm 0.05$	GAL
Sep. 01.35 1998	VLT-UT1	1500	$R_c > 25.29 \pm 0.06$	GAL

TABLE 2

B, V, R_c , I_c magnitudes, associated random errors ΔB , ΔV , ΔR_c , ΔI_c , and the coordinates of the reference stars. We estimate the absolute calibration uncertainty to be 0.05 magnitudes in B, V, and R_c and 0.10 magnitudes in I_c .

Star		A	B	C	D	E	F	G	H
B		21.693	19.614	23.161	23.654	22.296	23.976		
ΔB		0.022	0.010	0.042	0.060	0.016	0.073		
V		20.345	18.793	21.782	21.972	20.648	22.137	23.237	23.003
ΔV		0.011	0.010	0.017	0.021	0.010	0.023	0.064	0.044
R_c		19.646	18.413	20.874	21.119	19.640	20.889	23.008	21.906
ΔR_c		0.040	0.040	0.025	0.032	0.010	0.035	0.103	0.067
I_c		18.787	17.758	20.099	20.501	18.665	19.279		20.923
ΔI_c		0.034	0.034	0.045	0.062	0.034	0.040		0.080
RA (J2000)	5:01:	48.9	49.3	47.8	48.7	45.9	45.6	47.4	46.4
Dec (J2000)	11:	45:59	45:43	46:56	47:01	47:12	47:24	46:56	46:54




FIG. 1.— The V-, R_c - and I_c -band lightcurves of GRB 970228 (magnitude versus time). R_c -band data obtained from interpolations between V and I_c are indicated with open symbols (see the discussion for details). Indicated are power-law fits, $F_\nu \propto t^\alpha$, to the detections (ignoring upper limits). Fits to all points are indicated by the dotted lines and fits to the late-time lightcurve (after 20 days) are indicated by the dashed lines (for the V band they overlap). Also indicated is a fit to the R_c -band light curve at intermediate times (between 3 and 50 days; thick line).




FIG. 2.— The optical/near-infrared spectral flux distribution of GRB 970228 at March 30.8, 1997 UT (• and upper-limit arrow); from left to right: K-band detection (Table 1) translated into an upper limit because it is consistent with detection of the host galaxy only (detection magnitude minus 3σ), J-band detection (Table 1) translated into an upper limit due to unknown J-band host magnitude (detection magnitude minus 3σ), V and I_c HST measurements from Table 1. Also shown is the spectral flux distribution of SN 1998bw redshifted to the redshift of GRB 970228 (○). Note that the spectral flux distribution is different from that of Reichart (1999) (See the discussion in Sect. 4).




FIG. 3.— The V-, R_c -, and I_c -band lightcurves of GRB 970228 (fluxes versus time). The dotted curves indicate power-law decays with $\alpha = -1.73$, and redshifted SN 1998bw light curves. The thick line is the resulting sum of SN and power-law decay light curves.